

STIC-ILL

426799/

NO 362386

From: Holleran, Anne
Sent: Tuesday, September 04, 2001 6:03 PM
To: STIC-ILL
Subject: refs. for 09/251,133

Examiner: Anne Holleran
Art Unit: 1642; Rm 8E03
Phone: 308-8892
Date needed by: ASAP

Please send me copies of the following :

1. Chien, J. et al. Mol. and Cell. Endocrinology (2001) 181(1-2): 69-79
2. Chien, J. et al. Int. J. of Cancer (2001) 91(1): 46-54
3. Chien, J. et al. Oncogene (1999) 18(22): 3376-3382
4. Wong, E.C.C. et al. Proc. Amer. Assoc. for Cancer Res. (1997) 38: 288
5. Rayford, W. et al. Prostate (1997) 30(3): 160-166
6. Xue-Zhang, Q. et al. Endocrine (1995) 3(6): 445-451
7. Shah, G.V. et al. Endocrinology (1994) 134(2): 596-602
8. Rayford, W. et al. J. of Urology (1994) 151(5 suppl): 490A
9. Rayford, W. et al. J. of Urology (1993) 149(4 suppl): 479A
10. Shah, G.V. et al. Prostate (N.Y.) (1992) 21(2): 87-97
11. Sagol, O. et al. Annals of Medical Sciences (1999) 8(1): 14-21
12. Sussenot, O. et al. Prostate (1998) 36(suppl. 8): 43-51
13. Hanna, F.W. et al. J. Endocrinol. (1997) 152(2): 275-281
14. Sim, S.J. et al. Annals of Clinical and Laboratory Science (1996) 26(6): 487-495
15. Watanabe, K. et al. Fukushima J. Medical Science (1995) 41(2): 141-152
16. Esik, O. et al. European J. Gynaecological Oncology (1994) 15(3): 211-216

8823994

BEST AVAILABLE COPY

TWO CASES OF DUODENAL GANGLIOCYTIC PARAGANGLIOMA: IMMUNOCYTOCHEMICAL CHARACTERISTICS

KAZUO WATANABE¹, HIROSHI HASEGAWA², HIDEO SAKUMA³,
MASAO TSUBOI⁴, SEIKI ITO⁵ and TOSHIMITSU SUZUKI¹

¹*Department of Pathology II, Fukushima Medical College, Fukushima*

²*Department of Oral Surgery, Fukushima Medical College, Fukushima*

³*Pathology Division, Oota General Hospital, Koriyama*

⁴*Internal Medicine, Kuwano Kyouritsu Hospital, Koriyama*

⁵*Internal Medicine, Niigata University School of Medicine*

(Received November 17, 1995)

Abstract Two cases of duodenal gangliocytic paraganglioma were studied by means of immunocytochemical methods using 41 kinds of antibodies. The tumors consisted of three histological types; carcinoid, ganglioneuroma and paraganglioma. Tumors of both cases exhibited immunoreactivity to at least one or as many as three of the following: calcitonin, calcitonin-gene related peptide, endocrine granule constituent, Leu7, neuropeptide Y and basic fibroblast growth factor. In addition, these tumors were also immunopositive for neuron specific enolase, S-100 protein, neurofilament protein, pancreatic polypeptide, chromogranin A, somatostatin, leu-enkephalin, substance P and vasoactive intestinal peptide, as has been described in previous reports. In one case, tumor cells were immunopositive for adrenocorticotropin, bombesin, gastrin releasing peptide, myelin basic protein, neuroendocrine marker and tyrosine hydroxylase. Moreover, paraganglioma cells of tumors showed both argyrophilia and argentaffinity. These results suggest that duodenal gangliocytic paraganglioma may originate from embryonic neuroinsular complex.

Key words: Gangliocytic paraganglioma, immunocytochemistry, neuroendocrine markers, histogenesis

INTRODUCTION

Gangliocytic paraganglioma is an uncommon, largely benign, small intestinal neoplasm that usually occurs in the second portion of the duodenum, rarely in the jejunum^{1,2)} and in the peripancreatic tissue adjacent to large vessels³⁾. This tumor is generally solitary but a case involving multiple tumors has been documented³⁾.

渡辺一男, 長谷川 博, 佐久間秀夫, 坪井正夫, 伊藤正毅, 鈴木利光

Reprint requests to: Kazuo Watanabe, Department of Pathology II, Fukushima Medical College, Fukushima City 960-12, Japan

Immunocytochemical characteristics of 52 cases in the literature²⁻¹²⁾ examined, with a focus on the common and uncommon neuroendocrine features of this tumor. Recently, we had the opportunity to study two cases of this tumor using light microscopy and the streptoavidin-biotin immunostaining technique. On the basis of the results obtained, a possible origin of this tumor is discussed.

CASE REPORT

Case 1: A 48-year-old man was hospitalized for epigastralgia. He had a past history of duodenal ulcer and had undergone gastroduodenectomy 7 years earlier. An X-ray study confirmed the presence of a smooth surfaced mass in the second portion of the duodenum. Endoscopic examination disclosed a tumor mass with an eroded surface of papilla Vateri. Surgical exploration permitted complete excision of the duodenal tumor by combined removal of the duodenum and pancreatic head. The postoperative course was uneventful.

Case 2: A submucosal tumor was found in the second portion of the duodenum of a 51-year-old man by x-ray and endoscopic examination. His chief complaint was epigastric pain. The mass was resected and the diagnosis of the frozen section was gangliocytic paraganglioma. Recovery was uneventful.

MATERIALS AND METHODS

Light Microscopy and Histochemistry

The surgical specimens were fixed in 10% buffered formalin and processed in the usual manner for light microscopic study. Tissue sections were stained with hematoxylin and eosin and were subjected to argyrophilic Grimelius and Fontana-Masson argentaffin analysis.

Immunocytochemistry

Immunocytochemical studies were carried out on the formalin-fixed, paraffin-embedded tissue by the method described by Hsu *et al.*¹³⁾, using a kit of biotin-labelled second antibody and streptoavidin coupled with peroxidase (Nichirei Corp., Tokyo, Japan). The antibodies and abbreviations used are listed in Table I. Anti-bombesin is bombesin specific without any cross reactivity with gastrin-releasing peptide (GRP), nor does anti-GRP antibody react with bombesin¹⁴⁾. The anti-human pancreatic polypeptide (hPP) and porcine PP antibodies used do not recognize neuropeptide Y (NPY) and anti-NPY antibody does not bind to hPP or PPP. This was confirmed by a preabsorption test of the anti-hPP or anti-PPP antibodies with NPY (1 μ g/ml, Sigma, St. Louis, MO). Every absorption test revealed positive staining of the tumor cells (data not shown). As the insulin antibody was raised in guinea pig, a guinea pig PAP (peroxidase-antiperoxidase complex) kit (IBL, Gunma, Japan) was used. The peroxidase reaction product was visualized by Graham-

Table 1. List of antibodies used

Category	Antibody to	Source	Working titer
Peptide	Adrenocorticotropin (ACTH)	(Dr. Ito)* ¹	1:200
	Bombesin	(Dr. Yanaihara)* ²	1:1,000
	Calcitonin	(Dr. Yamada)* ¹	1:1,000
	Calcitonin-gene related peptide (CGRP)	(Peninsula)	1:3,000
	Chromogranin A	(Dakopatts)	1:500
	Endocrine granule constituent (EGC)	(MILAB)	1:500
	α -Endorphin	(Dr. Ito)	1:500
	β -Endorphin	(Dr. Ito)	1:500
	leu-Enkephalin	(Dr. Ito)	1:500
	Gastrin	(Dakopatts)	1:5,000
	Glucagon-like immunoreactivity (GLI)	(IBL)	1:500
	Glucagon	(IBL)	1:500
	Gastrin releasing peptide (GRP)	(Dr. Yanaihara)	1:2,000
	Insulin	(Dr. Ito)	1:500
	Neuroendocrine marker	(Lipshaw)	1:1
	Neuropeptide Y (NPY)	(Dr. Ito)	1:500
	Neurotensin	(Dr. Yanaihara)	1:1,000
	Human pancreatic polypeptide (hPP)	(Dr. Ito)	1:200
	Porcine pancreatic polypeptide (PPP)	(Dr. Yanaihara)	1:1,000
	Parathormone	(IBL)	1:500
	Somatostatin	(Dr. Ito)	1:100
	Substance P	(Dr. Yanaihara)	1:1,000
	Vasoactive intestinal polypeptide (VIP)	(Dr. Yanaihara)	1:500
Neuronal marker	Brain natriuretic peptide (BNP)	(IBL)	1:500
	Glial fibrillar acidic protein (GFAP)	(IBL)	1:1,000
	Myelin basic protein (MBP)	(IBL)	1:500
	Neurofilament protein (NFP)	(Boehringer)	
	68 kDa		1:4
	160 kDa		1:4
	200 kDa		1:4
Growth factor	Neuronspecific enolase (NSE)	(IBL)	1:500
	S-100 protein	(IBL)	1:500
	Acidic fibroblast growth factor (a-FGF)	(Dr. Sasaki)* ³	1:500
	Basic fibroblast growth factor (b-FGF)	(Dr. Sasaki)	1:500
	Insulin-like growth factor-II (IGF-II)	(Cosmo-Bio)	0.1 μ g/ml
Others	Carcinoembryonic antigen (CEA)	(IBL)	1:1,000
	Epithelial keratin (AE-1, AE-3)	(ICN Immunobiol.)	1:100
	Leu-7	(Becton Dickinson)	0.2 μ g/ml
	Serotonin	(Dr. Yui)* ¹	1:1,000
	Tyrosine hydroxylase	(Dr. Kumanishi)* ¹	1:400

*¹ Niigata University, Niigata, Japan.*² University of Shizuoka, Shizuoka, Japan*³ Bio

Karnovsky solution¹⁵⁾ containing 20 mg/dl of 3, 3'-diaminobenzidin tetrahydrochloride (Dojin, Kumamoto, Japan) and nuclei were stained by 1% methyl green. Endogenous peroxidase activity was blocked by 0.3% H₂O₂ in methanol prior to immunostaining and also by 65 mg/dl of sodium azide (Wako Pure Chem. Indust., Osaka, Japan) in Graham-Karnovsky solution.

RESULTS

Light microscopy

Tumors of both patients were well-circumscribed in the duodenal wall, as shown in Fig. 1a, and involved the submucosa, with a focal ulceration in case 1. Three histological patterns were observed in both tumors; paraganglioma, ganglioneur-




Fig. 1. In case 1, the tumor in the duodenal wall is well-circumscribed and involves the submucosa with focal ulceration of the overlying mucosa (arrowhead, a). Both tumors show three histological patterns: paraganglioma (b), ganglioneuroma (c) and carcinoid (d). a; $\times 5$, b-d; $\times 100$.

oma and carcinoid. In case 1, ganglioneuroma was predominant, whereas in case 2, paraganglioma and ganglioneuroma were equally observed. The carcinoid type was the least represented in both cases. Amyloid, as described in the previous report by Reed *et al.*¹¹⁾, was not present.

The paraganglioma involved the lamina propriae of the duodenal mucosa in both cases and was characterized by a rounded nest of epithelioid cells demarcated

Table 2. Histo- and Immunocyto-chemical Profile of Gangliocytic Paraganglioma

Category	Case		Histological component positively stained
	1	2	
Histochemistry			
Grimelius	+	+	paraganglioma
Masson-Fontana	+	+	paraganglioma
Peptide			
Adrenocorticotropin	—	+	paraganglioma
Bombesin	—	+	paraganglioma
Calcitonin	+	+	carcinoid/ganglioneuroma
Calcitonin gene related peptide	‡	‡	carcinoid/ganglioneuroma/paraganglioma
Chromogranin A	‡	‡	carcinoid/ganglioneuroma/paraganglioma
Endocrine granule constituent	‡	‡	carcinoid/ganglioneuroma/garaganglioma
leu-Enkephalin	‡	‡	ganglioneuroma/paraganglioma
Gastrin releasing peptide	—	‡	carcinoid/paraganglioma
Neuroendocrime marker	—	+	paraganglioma
Neuropeptide Y	‡	+	paraganglioma
Pancreatic polypeptide, human	‡	+	carcinoid/paraganglioma
Pancreatic polypeptide, porcine	‡	+	carcinoid/ganglioneuroma/paraganglioma
Somatostatin	‡	‡	carcinoid/ganglioneuroma/paraganglioma
Substance P	—	+	paraganglioma
Vasoactive intestinal peptide	+	+	ganglioneuroma/paraganglioma
Nouronal marker			
Myelin basin protein	—	+	ganglioneuroma
Neurofilament protein			
160kDa	‡	‡	ganglioneuroma
200kDa	‡	‡	ganglioneuroma
Nouron specific enolase	‡	‡	carcinoid/ganglioneuroma/paraganglioma
S-100 protein	‡	‡	ganglioneuroma
Growth factor			
basic fibroblast growth factor	+	+	ganglioneuroma/paraganglioma
Others			
Leu7	+	‡	carcinoid/ganglioneuroma/paraganglioma
Tyrosine hydroxylase	—	+	ganglioneuroma/paraganglioma

Score: + ; slightly immunoreactive, # ; highly immunoreactive

- ; no immunoreactivity

by thin, delicate, vascularized fibrous septa (Fig. 1b). The ganglioneuroma was composed of clusters of ganglion cells in which nuclei were frequently pyknotic and contained fibrous connective tissue matrix (Fig. 1c). The carcinoid consisted of areas in which neoplastic cells had formed ribbons, festoons and small glands in a manner indistinguishable from the usual type of carcinoid (Fig. 1d). Some of the carcinoid tumor cells and paraganglioma cells had intimately intermingled.

Histochemistry and immunocytochemistry

Table 2 shows the histochemical and immunocytochemical reactivities of the neoplasm. Reactivities of the latter type were scored as indicated in the footnotes to the Table 2. Both argentaffin and argyrophilic granules were observed in the cytoplasm of paraganglioma cells in both tumors. Positive cells, although not so numerous, were polygonal or fibrocytic and were located at the periphery of the




Fig. 2. Immunolabelled tumor cells with antibodies against ACTH (a), bombesin (b), CT (c) and CGRP (d) in the paraganglioma. a-d; $\times 200$.




Fig. 3. Carcinoid tumor cells positively immunostained for human PP (a), and GRP (b); paraganglioma tumor cell stained positive for NPY (c) and EGC (d). a, c, d; $\times 200$, b; $\times 100$.

tumor nests (not shown). Positive immunoreactivity was detected to 12 peptides, including adrenocorticotropin (ACTH), bombesin, calcitonin (CT), calcitonin gene-related peptide (CGRP) (Fig. 2a-d), HPP, GRP, NPY (Fig. 3a-c), leu-enkephalin, PPP, somatostatin, substance P and vasoactive intestinal polypeptide (VIP). Positivity to neuroendocrine cell markers, such as chromogranin A, endocrine granule constituent (EGC) (Fig. 3d), Leu7 (Fig. 4a), tyrosine hydroxylase (Fig. 4b) and neuroendocrine marker, was found as well. Other proteins detected in the tumors included basic fibroblast growth factor (bFGF), although its reactivity was weak, myelin basic protein (MBP), 200 kDa (not shown) and 160kDa neurofilament protein (Fig. 4d), neuron specific enolase (NSE) and S-100 protein. The following immunoreactivities were also noted but only in case 2; these were GRP in paraganglioma and carcinoid, ACTH and bombesin in paraganglioma, MBP in fibers of ganglioneuroma, neuroendocrine marker in paraganglioma, and tyrosine hydroxylase in paragangli-




Fig. 4. Positive immunocytochemical staining of Leu7 in the ganglioneuroma (a), tyrosine hydroxylase in the paraganglioma (b), basic FGF in the paraganglioma (c) and NFP (160kDa) in the ganglioneuroma (d). a, b; $\times 100$, c; $\times 200$, d; $\times 40$.

oma. Among the immunoreactivities tested, CGRP, NFPs, Leu7, NSE, PPs, somatostatin and S-100 protein were frequently observed.

DISCUSSION

Gangliocytic paraganglioma was first reported by Dahl *et al.* in 1957¹⁶⁾. A few years later in 1962, Taylor and Helwig described nine cases of nonchromaffin paraganglioma of the duodenum¹⁷⁾. Since then, about 96 additional cases have been reported^{1,3-11,18-29)}. Almost all of these cases demonstrated argyrophilia but not argentaffinity. Our two cases represent the first reported instances of both argyrophilia and argentaffinity, albeit within a limited area. Although the argentaffin method can be used to visualize serotonin storing cells, serotonin immunoreactivity was not present in our cases. Therefore, the significance of this argentaffin-positive

reaction is not clear, however, the possible origin of this tumor will be discussed later. Fifty-two reported cases were examined with respect to their immunohistochemical features, as stated earlier. Results^{2,4,5,7,9,29)} revealed that NSE, S-100 protein, NFP, PP, chromogranin A, somatostatin, leu-enkephalin, and protein gene product 9.5 were frequently detected in tumor tissue, whereas CCK, molluscan cardioexcitatory peptide, serotonin, gastrin, VIP, glucagon, insulin and substance P were only infrequently mentioned^{11,12)}. In the current study, we found many tumor cells which showed positive immunoreactivity for PP, somatostatin, leu-enkephalin, NSE, S-100 protein, NFPs, and chromogranin A as was the cases in the previous reports. In addition, CGRP, EGC, Leu7 and NPY immunoreactive tumor cells were found frequently in our two cases. Moreover, calcitonin, b-FGF, MBP and TH immunoreactivities were detected as well, although the frequencies were rather low.

The overall immunocytochemical features of the tumors presented seem to be similar to those of pheochromocytoma³⁰⁻³⁴⁾ and paraganglioma³⁵⁾, but the frequent detection of PP-like immunoreactivity in this tumor is a distinctive finding, since the usual form of paraganglioma or pheochromocytoma does not show this PP-like immunoreactivity. Further discussion related to this will appear later.

The origin of this tumor is still a matter of controversy because of histological complexity. However, on the basis of morphological and immunohistochemical analyses, several sources have been implicated. These include pluripotent stem cells residing at the base of intestinal glands³⁾, undifferentiated neural crest cells³⁹⁾, multipotential cells of the embryonic celiac ganglion¹⁷⁾, residual paraganglionic cells and Meissner's plexus³⁶⁾, a paraganglionic vagal branch in the duodenum³⁷⁾, and embryonic neuroinsular complex⁵⁾ which was first described in the pancreas of sheep embryo by van Campenhout³⁸⁾. The first three of the postulated sources are unlikely, because pluripotent cells capable of differentiating to endodermal endocrine cells and neuroepithelial ganglionic cells have not been found in the duodenum. Residual paraganglionic cells and Meissner's plexus both seem to be an unlikely origin since paraganglionic tissue has not been described in the duodenal wall and the authors did not speculate why Meissner's plexus and residual paraganglion cells would act in concert to produce a neoplasm³⁶⁾. The last speculation seem to be rather feasible since sympathetico-insular complex does exist not only in the pancreas but also in the small bowel and appendix of mammalian embryos^{39,40)}. These endodermal-neuroectodermal complexes are thought to be composed of argentaffin cells that migrated from the crypts of Lieberkühn into the lamina propriae and submucosa, where they maintain intimate contact with nerve fibers and ganglion cells⁴⁰⁾. This finding explains the argentaffinity of the cells of this tumor. In human embryo of five-months gestation, Osaka and Kobayashi⁴¹⁾ described basigranulated endocrine cells intermingled with Schwann-like cells in the lamina propriae of duodenum and termed this association neuro-endocrine complex. This structure is thought to be homologous to the "complexe sympathico-insulaire"³⁸⁾, later revised to the "complexe neuroinsulaire"³⁹⁾, in mammalian embryo, and also to

the neuroendocrine complexes in the avian⁴²⁾ and rat stomach⁴³⁾. From these observations it is strongly suggested that this peculiar gangliocytic paraganglioma might be derived from these complexes exhibiting carcinoid and ganglioneuroma/paraganglioma patterns.

The frequent occurrence of the PP-like immunoreactive tumor cells in the duodenal paragangliocytic paraganglioma might not be a distinctive feature of this tumor since some duodenal carcinoid tumors have been reported to contain PP-like tumor cells⁴⁴⁾. In our tumors, PP-like immunoreactivity was found not only in the carcinoid but also in the paraganglioma. This suggests a possible transition of carcinoid tumor cells to paraganglioma-like cells, since the carcinoid cells in our cases were intimately entangled with paraganglioma cells suggesting this type of transition. More directly, the transformation of endocrine (carcinoid) tumor cells into neuron-like cells have been observed *in vitro*⁴⁵⁾. These authors reported that tumor cells spontaneously transform into cells with a neuron-like morphology and express neurofilament protein and synaptic vesicle membrane component.

In conclusion, our study indicates that duodenal gangliocytic paraganglioma exhibits many types of neuroendocrine immunoreactivity which are also shared with ganglioneuromas and paragangliomas occurring in sites other than the duodenum as well as with duodenal carcinoids.

REFERENCES

1. Reed RJ, Caroca PJ, Harkin JC, Gangliocytic paraganglioma. *Am J Surg Pathol*, 1, 207-216, 1977.
2. Burke AP, Helwig EB, Gangliocytic paraganglioma. *Am J Clin Pathol*, 92, 1-9, 1989.
3. Kawaguchi K, Takizawa T, Koike M, Tabata I, Goseki N, Multiple paraganglioneuromas. *Virchows Arch A Pathol Anat*, 406, 373-380, 1985.
4. Guarda LA, Ordonez, NZ, DelJunco GW, Luna MA, Gangliocytic paraganglioma of the duodenum: An immunocytochemical study. *Am J Gastroenterol*, 78, 794-798, 1983.
5. Perrone T, Sibley RK, Rosai J, Duodenal gangliocytic paraganglioma, an immunohistochemical and ultrastructural study and a hypothesis concerning its origin. *Am J Surg Pathol*, 9, 31-41, 1985.
6. Hamid QA, Bishop AE, Rode J, Dhillon AP, Rosenberg BF, Reed RJ, Sibley RK, Polak JM, Duodenal gangliocytic paragangliomas: A study of 10 cases with immunocytochemical neuroendocrine markers. *Hum Pathol*, 17, 1151-1157, 1986.
7. Anders KH, Glasgow BJ, Lewin KJ, Gangliocytic paraganglioma associated with duodenal adenocarcinoma. Case report with immunohistochemical evaluation. *Arch Pathol Lab Med*, 111, 49-52, 1987.
8. Stephans M, Williams GT, Jasani B, Williams ED, Synchronous duodenal neuroendocrine tumors in von Recklinghausen's disease- a case report of co-existing gangliocytic paraganglioma and somatostatin-rich glandular carcinoid. *Histopathology*, 11, 1331-1340, 1987.
9. Barbareschi M, Frigo B, Aldovini D, Leonardi E, Cristina S, Falleni M, Duodenal gangliocytic paraganglioma. Report of a case and review of the literature. *Virchows Arch A Pathol Anat*, 416, 81-89, 1989.
10. Grouls V, Stein U, Vogel J, Juxtapapillares gangliozytisches Paragangliom des Duodenums. *Dtsch Med Wschr*, 114, 584-588, 1989.

11. Barten M, Kloppel G, Das gangliozytische Paragangliom des Duodenums. Fallbericht mit immunozytochemischer Charakterisierung. *Zentralb Allg Pathol Pathol Anat*, 136, 181-187, 1990.
12. Collina G, Maiorana A, Trentini GP, Duodenal gangliocytic paraganglioma. Case report with immunohistochemical study on the expression of keratin polypeptides. *Histopathology*, 19, 476-478, 1991.
13. Hsu S-M, Raine L, Fanger H, Use of avidin-biotin-peroxidase complex (ABC) in immunoperoxidase techniques. A comparison between ABC and unlabelled antibody (PAP) procedures. *J Histochem Cytochem*, 29, 577-580, 1981.
14. Matubayashi S, Yanaihara C, Ohkubo M, Fukata S, Hayashi Y, Tamai H, Nakagawa T, Miyauchi A, Kuma K, Abe K, Suzuki T, Yanaihara N, Gastrin-releasing peptide immunoreactivity in medullary thyroid carcinoma. *Cancer*, 53, 2472-2477, 1984.
15. Graham RC, Karnovsky MJ, The early stage of absorption of injected horseradish peroxidase in proximal tubules of mouse kidney: Ultrastructural cytochemistry by a new technique. *J Histochem Cytochem*, 14, 291-302, 1966.
16. Dahl EV, Waugh JM, Dahlin DC, Gastrointestinal ganglioneuromas. Brief review with report of a duodenal ganglioneuroma. *Am J Pathol*, 33, 953-966, 1957.
17. Taylor HB, Helwig EB, Benign nonchromaffin paragangliomas of the duodenum. *Virchows Arch A Pathol Anat*, 335, 356-366, 1962.
18. Lukash WM, Hyams VJ, Nielsen OF, Neurogenic neoplasms of the small bowel: Benign non-chromaffin paraganglioma of the duodenum. Report of a case. *Am J Dig Dis*, 11, 575-579, 1966.
19. Germer M, Feutchwanger MM, Ganglioneuroma of the duodenum. *Gastroenterology*, 51, 689-693, 1966.
20. Goldman RL, Ganglioneuroma of the duodenum. Relationship to non-chromaffin paraganglioma of the duodenum. *Am J Surg*, 115, 716-719, 1968.
21. Weitzner S, Benign nonchromaffin paraganglioma of the duodenum. Report of a case and review of the literature. *Am J Gastroenterology*, 53, 365-369, 1970.
22. Kepes JJ, Zacharias DL, Gangliocytic paraganglioma of the duodenum. A report of two cases with light and electron microscopic examination. *Cancer*, 27, 67-70, 1971.
23. Lauzon A, Cadotte M, Paragangliome gangliocytaires du duodenum. *L'Union Med Can*, 101, 1584-1586, 1972.
24. Qizilbash AH, Benign paraganglioma of the duodenum. A case report with light and electron microscopic examination. *Arch Pathol*, 96, 276-280, 1973.
25. Kermarec JJ, Duplay H, Lesbron F, Paragangliome gangliocytique du duodenum. Une observation avec etude ultra-structurale. *Arch Anat Cytol Pathol*, 24, 261-268, 1976.
26. Cooney T, Sweeney EC, Paraganglioma of the duodenum: an evolutionary hybrid? *J Clin Pathol*, 31, 233-244, 1978.
27. Babaryka I, Polypoid nonchromaffin paraganglioma of the duodenum. *Virchows Arch A Pathol Anat*, 377, 181-187, 1978.
28. Beltrami CA, Montironi R, Cinti S, Gangliocytic paraganglioma of the duodenum: case report. *Tumori*, 66, 637-641, 1980.
29. Cohen T, Zweig SJ, Tallis A, Auazon R, Reich M, Paraganglioneuroma of the duodenum. Report of a case with radiographic findings, angiographic findings and review of the literature. *Am J Gastroenterology*, 75, 197-203, 1981.
30. Adrian TE, Allen JM, Terenghi G, Bacarese-Hamilton AJ, Brown MJ, Polak JM, Neuropeptide Y in pheochromocytomas and ganglioneuroblastomas. *Lancet* ii, 540-542, 1983.
31. Tsutsumi Y, Leu7 immunoreactivity as histochemical marker for paraffin-embedded neuroendocrine tumors. *Acta Histochem Cytochem*, 17, 15-21, 1984.
32. Osamura RY, Tsutsumi Y, Yanaihara N, Imura H, Watanabe K, Immunohistochemical studies for multiple peptide-immunoreactivities and co-localization of met-enkephalin-Arg⁶-Gly⁷-Leu⁸, neuropeptide Y and somatostatin in human adrenal medulla and pheo-

- chromocytomas. *Peptides*, 8, 77-87, 1987.
33. Hachker GW, Bishop AE, Terenghi G, Varndell IM, Aghahowa J, Pollard K, Thurner J, Polak JM, Multiple peptide production and presence of general neuroendocrine markers detected in 12 cases of human pheochromocytoma and in mammalian adrenal glands. *Virchows Archiv A Pathol Anat*, 412, 399-411, 1988.
 34. Kimura N, Functioning and nonfunctioning adrenal medullary tumors. *Endocrine Pathol*, 2, 64-73, 1991.
 35. Takahashi H, Nakashima S, Kumanishi T, Ikuta F, Paraganglioma of the craniocervical region. An immunohistochemical study on tyrosine hydroxylase. *Acta Neuropathol*, 73, 227-232, 1987.
 36. Konok GP, Sanchez-Cassis G, Gangliocytic paraganglioma of the duodenum. *Canad J Surg*, 22, 173-175, 1979.
 37. Böck P, The paraganglia. Springer-Verlag, Berlin Heidelberg, and New York, 1982.
 38. van Campenhout E, Contribution a l'etude de l'histogenese du pancreas chez quelques mammiferes. Les complexes sympathico-insulaires. *Arch Biol (Liege)*, 37, 121-171, 1927.
 39. van Campenhout E, Contribution au probleme des connexions neuro-entoblastiques. *Acad R Med Belg*, 5, 189-201, 1940.
 40. van Campenhout E, Les cellules argentaffines de l'appendice ileo-caecal de l'embryon humain. *Acad R Med Belg*, 18, 160-182, 1953.
 41. Osaka M, Kobayashi S, Duodenal basal-granulated cells in the human fetus with special reference to their relationship to nervous elements. In "Endocrine Gut and Pancreas" (ed. by Fujita, T.) Elsevier, Amsterdam, pp. 145-158, 1976.
 42. Kataoka K, An electron microscopic study on a neuroendocrine complex in the proventricular mucosa of the finch, *Uroloncha striata* var *domestica*. *Arch Histol Jap*, 36, 391-400, 1974.
 43. Matsuo Y, Seki A, Fukuda S, Neuroendocrine cells in the lamina propria of rat stomach. In "Endocrine Gut and Pancreas" (ed. by Fujita, T.) Elsevier, Amsterdam, pp. 159-165, 1976.
 44. Iwafuchi M, Watanabe H, Noda Y, Ajioka Y, Enjoji M, Ito S, Gastrointestinal carcinoid tumors of Japanese: incidence and characteristics based on anatomical classification, with special reference to difference between carcinoid tumor and endocrine cell carcinoma. *I to Cho*, 24, 869-882, 1989. (in Japanese)
 45. Ahlman H, Ahlund L, Dahlstrom A, Teodorsson E, Expression of gastrointestinal endocrine tumors in culture system. *Arch Histol Cytol*, 52(Suppl.), 233-240, 1989.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☐ FADED TEXT OR DRAWING

☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.